

Wildfire Smoke in the United States

Jacob Gellman* and Matthew Wibbenmeyer†

Introduction

Wildfire activity is increasing globally (Jolly et al. 2015; Senande-Rivera, Insua-Costa, and Miguez-Macho 2022), driven by increases in aridity and, in some regions, an overabundance of fire-ready fuels (Marlon et al. 2012). Recently in the United States, fires have produced several deadly and destructive disasters, including those in Paradise, California, in 2018, and Lahaina, Hawaii, in 2023. Despite these notable impacts, there is growing evidence that the overall damages of wildfire smoke emissions may be at least as large as the direct damages from wildfires (Wang et al. 2021; Burke et al. 2023).

As fire consumes biomass, it releases a variety of pollutants, including carbon dioxide, carbon monoxide, methane, volatile organic compounds, nitrous oxide, nitrogen oxides, and particulate matter. Of these, fine particulate matter less than 2.5 microns in diameter ($PM_{2.5}$) accounts for the third-largest share of emissions, by mass, after carbon dioxide and carbon monoxide (Urbanski, Hao, and Baker 2008).¹ Researchers have found that even short-term exposure to heightened levels of fine particulate pollution can have substantial health impacts because of its effects on cardiovascular and respiratory systems. Heightened $PM_{2.5}$ levels can lead to increased mortality, especially among vulnerable individuals, and to increased hospitalizations and preterm births. $PM_{2.5}$ can also reduce welfare through other channels; for example, it has been found to decrease economic productivity, increase crime rates, and impair educational outcomes.

This article argues that increasing air pollution due to wildfire smoke is a serious threat to health and human well-being and warrants increased attention from policy makers in the United States. Although acknowledging the important role of carbon emissions from wildfires in contributing to climate change—for example, in 2020, only California's transportation sector produced more emissions than the state's wildfires (Jerrett, Jina, and Marlier 2022)—this

*Department of Economics, University of Alaska Anchorage; †Resources for the Future (corresponding author; email: wibbenmeyer@rff.org)

¹Some literature has also studied exposure to PM_1 and PM_{10} , which are particulate matter less than 1 and 10 μm in diameter, respectively. Studies generally indicate that exposure to particulate matter less than 2.5 microns in diameter is especially harmful. Carbon monoxide and carbon dioxide account for 92–95 percent and 4–7 percent of the composition of wildfire smoke emissions, respectively, depending on forest type; $PM_{2.5}$, which is the third-largest constituent emission, accounts for 0.09–0.8 percent of emissions (Urbanski, Hao, and Baker 2008).

Electronically published January 10, 2025

Review of Environmental Economics and Policy, volume 19, number 1, winter 2025.

© 2025 Association of Environmental and Resource Economists. All rights reserved. Published by The University of Chicago Press for AERE. <https://doi.org/10.1086/732404>

article focuses specifically on wildfire smoke, that is, emissions that contribute conventional pollutants. Agricultural burning and prescribed fire, another category of wildland fire, can also contribute to air pollution; however, we restrict our attention to smoke from wildfires—uncontrolled fires burning on wildlands such as forests and shrublands.

The article proceeds as follows. “Impacts of Wildfire Smoke” reviews evidence from epidemiologic and economic studies on the impacts of wildfire smoke, mainly focusing on studies that have specifically investigated wildfire smoke but occasionally drawing from a large body of evidence on the effects of $PM_{2.5}$ more broadly. “Wildfire Smoke Trends” then shows, based on recently published results and data, that the impact of wildfire smoke on US air quality has grown substantially in recent years. Finally, “Policy Responses” outlines potential paths for policy to address increasing smoke and discusses the considerations surrounding smoke from prescribed fires, which reduce future wildfire smoke impacts but produce pollution in the immediate term. The final section concludes.

Impacts of Wildfire Smoke

Health

Most studies of the effects of wildfire smoke on health have focused on harmful effects due to particulate pollution. Among air pollutants, fine particulate matter is one of the most harmful to human health. Its small size allows it to easily enter the lungs and eventually escape into the bloodstream, with deleterious effects throughout the body. $PM_{2.5}$ particles cause inflammation in the airway, lungs, and central nervous system, and they change cell chemistry, resulting in decreased function of the respiratory and cardiovascular system (Feng et al. 2016). These effects can exacerbate underlying conditions such as chronic pulmonary disease and asthma and increase the risk of heart attack, congestive heart failure, and dementia, especially among the elderly. $PM_{2.5}$ may also increase the incidence of preterm birth, low birth weight, and infant mortality (for reviews of the physiological mechanisms underlying health effects of fine particulate matter, see Kim, Kabir, and Kabir 2015 and Feng et al. 2016).

Economists have studied the impacts of $PM_{2.5}$ from wildfire smoke because it generates arguably exogenous temporal variation in fine particulate concentrations, allowing them to econometrically separate the health effects of $PM_{2.5}$ from the effects of variables correlated with both pollution exposure and health, such as race and income. Researchers frequently rely on satellite data to measure the presence of wildfire smoke (e.g., Ruminiski et al. 2008). These data do not distinguish between ground-level smoke and smoke at higher altitudes, however, so it is important to couple satellite data with surface-level air quality measures (e.g., Burkhardt et al. 2019; Burke et al. 2022).

The health consequences of exposure to $PM_{2.5}$ from wildfire smoke versus other sources may differ for two reasons. First, $PM_{2.5}$ from wildland fires typically exhibits considerably more temporal variation than $PM_{2.5}$ from the industrial and transportation sectors. Exposure from wildfire smoke is more likely to be acute and is often more severe; for example, during June 2023, New York City’s air quality was briefly the worst in the world because of smoke from Canadian wildfires. Second, although $PM_{2.5}$ is defined by diameter, the chemical composition of particulates can vary, and $PM_{2.5}$ emissions from wildfires differ chemically from

those of other sources (Wegesser, Pinkerton, and Last 2009). In addition, smoke contains harmful co-pollutants, including larger-diameter particulate matter and volatile organic compounds (Liu and Peng 2019). Aguilera et al. (2021) found that wildfire smoke increases respiratory hospitalizations more than $PM_{2.5}$ from ambient sources.

In general, however, epidemiological and econometric studies find health effects from smoke that are broadly consistent with the greater literature on health effects of *acute* $PM_{2.5}$ exposure. There is strong evidence that wildfire smoke increases mortality (Reid et al. 2016; Cascio 2018). Qiu et al. (2024) estimate that from 2011 to 2020 there were approximately 15,800 excess deaths per year due to wildfire smoke. Johnston et al. (2012) and Roberts and Wooster (2021) each estimate that, globally, landscape fires (including both wildland fires and agricultural burning) cause as many as 600,000 additional deaths per year. However, more research is needed to clarify the magnitude and primary mechanisms that underlie mortality effects (Liu et al. 2015; Reid et al. 2016; Cascio 2018). Although there is consistent evidence that wildfire smoke increases hospital visits for respiratory conditions (Reid et al. 2016; Heft-Neal et al. 2023), evidence of smoke's effects on cardiovascular morbidity is mixed (Heft-Neal et al. 2023).

Studies have also shown that wildfire smoke affects birth outcomes. For example, in California, Heft-Neal et al. (2022) estimate that each additional day of exposure to wildfire smoke during pregnancy is associated with a 0.49 percentage point increase in the probability of preterm birth. Similarly, using a difference-in-differences approach, McCoy and Zhao (2021) show that exposure to wildfire smoke causes a 3.4 percentage point increase in the probability of low birth weight.

Welfare

In addition to direct health costs, smoke indirectly affects economic welfare through a variety of channels. Sickness due to smoke, or efforts to avoid sickness, can result in reduced work hours and earnings. Smoke can impair cognitive function, which may affect productivity, educational outcomes, and crime. The overall perceived welfare impacts of these various effects can be measured based on survey responses or preferences revealed through individual behavior, such as demand for housing.

Borgschulte, Molitor, and Zou (2024) find that wildfire smoke reduces quarterly earnings in the United States by an average of \$125 billion per year—similar to mortality losses due to smoke. They attribute approximately 13 percent of observed earnings losses to reduced employment, with the remaining losses presumably from reduced hours or wages. Several studies in the air pollution literature have similarly documented a reduction in working hours due to $PM_{2.5}$ (Aragon, Miranda, and Oliva 2017; Fan and Grainger 2023).

In addition to reduced hours, particulate matter may impair labor productivity. Although evidence on the effects of particulate matter on productivity is mixed (Chang et al. 2016, 2019; He, Liu, and Salvo 2019), there is substantial evidence that particulate pollution negatively affects the *inputs* into productivity. $PM_{2.5}$ has been found to cause declines in performance on various cognitive tasks (Graff Zivin and Neidell 2012; Archsmith, Heyes, and Saberian 2018; Bedi et al. 2021; Schmidt 2022). Particulate pollution, including that from wildfire smoke, negatively affects school performance and attendance (Chen, Guo, and Huang 2018; Wen and Burke 2022), which may have longer-run effects on productivity. Psychological effects of particulate pollution include increased stress and anxiety (Power et al. 2015; Sass

et al. 2017). Burkhardt et al. (2019) propose these psychological phenomena as the primary mechanism to explain estimated increases in violent crime due to particulate pollution.

Smoke also has direct effects on agricultural output. Low-density smoke plumes increase the proportion of diffuse light, which can increase crop yields; because wildfires are currently more likely to produce low-density plumes, the net effect is to increase crop yields, but these benefits are expected to dissipate by 2050 as wildfires become more frequent and severe (Behrer and Wang 2022). Nevertheless, smoke can be detrimental to particular crops, such as wine grapes (Whiting and Krstic 2007).

An alternative to studying smoke damages using observational data on market outcomes is to measure the value of damages using survey responses or observed individual behavior. These approaches could potentially undervalue smoke impacts—for example, if individuals do not fully appreciate the health consequences of exposure. An advantage of both survey and observational data, however, is that they can account for effects on the subjective value of amenities (such as smoke-free air) or well-being. For example, Jones (2017) compares life satisfaction survey responses across smoky and nonsmoky periods and finds that US adults are willing to pay \$373 to avoid one day of wildfire smoke. Similarly, Burke et al. (2022) and Du et al. (2024) estimate effects of wildfire smoke on expressed sentiment using high-frequency social media data, with Du et al. (2024) translating these effects to reductions in wages in Southeast Asia. Gellman, Walls, and Wibbenmeyer (2023) examine welfare effects using observed recreation behavior, finding that smoke causes welfare losses to outdoor recreation of approximately \$2.3 billion per year in the western United States.

Finally, preferences regarding smoke may be visible in home prices and regional demand for housing. Several studies find that air pollution more generally affects willingness to pay for housing (Chay and Greenstone 2005; Hamilton and Phaneuf 2015; Freeman et al. 2019; Nam et al. 2022). In general, it may be difficult to disentangle the effects of smoke from regional shocks to home prices. However, it is reasonable to expect that repeated smoke events might affect housing demand, and some initial evidence indicates that it does (Huang and Skidmore 2024). Because smoke tends to be spatially correlated, reduced demand for homes may result in increased outmigration from heavily affected regions (Chen, Oliva, and Zhang 2022). Although no studies have empirically examined migration due to wildfire smoke specifically, Rubin and Wong-Parodi (2022) find in a survey of California residents that nearly a quarter of those who intended to move within the next five years reported that wildfire and smoke had at least a moderate effect on their migration decision.

Distributional Impacts

Welfare effects from smoke—through either health impacts or other channels—are not borne equally. Although it is well established that disadvantaged and non-White communities are more exposed to overall $PM_{2.5}$, counties with a higher proportion of White people are on average more exposed to $PM_{2.5}$ from wildfire smoke (Burke et al. 2021). However, conditional on exposure, the degree of vulnerability to smoke varies across populations because of differences in impacts on indoor air quality, protective behavior, and time spent outdoors (Liang et al. 2021; Marlier et al. 2022; O'Dell et al. 2022; Wen and Burke 2022).

On average, an increase in outdoor $PM_{2.5}$ of one microgram per cubic meter ($\mu\text{g}/\text{m}^3$) is associated with an increase in indoor $PM_{2.5}$ of 0.15–0.4 $\mu\text{g}/\text{m}^3$. Indoor air pollution can triple

during smoke events, in some cases leading to indoor $PM_{2.5}$ concentrations that exceed the $35 \mu\text{g}/\text{m}^3$ standard for daily ambient air quality set by the EPA (Liang et al. 2021; Burke et al. 2022). However, infiltration varies by building type (Liang et al. 2021; Burke et al. 2022; O'Dell et al. 2022) and is higher for older homes, low-income neighborhoods, and Black and Hispanic neighborhoods (Burke et al. 2022).

Outdoor workers are more exposed than indoor workers to hazardous air quality. Outdoor workers constitute 10 percent of the US workforce (BLS 2019) and earn substantially less than indoor workers, with the mean at \$46,400 per year for outdoor workers and \$54,300 per year for indoor workers.² Hispanic individuals are overrepresented among outdoor workers (Cox-Ganser and Henneberger 2021).

The ability to engage in protective behavior also varies across socioeconomic populations (Burke et al. 2022; O'Dell et al. 2022). Defensive behaviors, including health-protecting investments like air purifiers (Richardson, Champ, and Loomis 2012; Ito and Zhang 2020), and avoidance behaviors, such as reducing short-run labor participation (Aragon, Miranda, and Oliva 2017; Borgschulte, Molitor, and Zou 2024), can both be costly. During smoke events, residents of lower-income areas search the internet less for health-protective information and spend less time at home than those in higher-income areas (Burke et al. 2022). These results show that adaptive behavior is likely to be unequal across populations, conditional on exposure.

Wildfire Smoke Trends

As a consequence of the increasing size and intensity of wildfires, pollution from wildfire smoke is rising. This increased smoke pollution is currently undermining federal air quality goals. Wildfire smoke has accounted for up to 25 percent of $PM_{2.5}$ in recent years across the United States and up to half in some areas of the western United States (O'Dell et al. 2019; Burke et al. 2021). Most exposure has been concentrated in the western United States, especially the Pacific Northwest (Burke et al. 2021, 2023). Although overall levels of ambient $PM_{2.5}$ pollution had been declining for several decades, wildfire smoke pollution has reversed those trends in 31 states (Burke et al. 2023). From 2011 to 2022, wildfire smoke accounted for at least 25 percent of days exceeding the EPA's 24-hour daily standards for $PM_{2.5}$ in seven states (Burke et al. 2023). These poor air quality days are generally not a result of fires burning within the same county; rather, 87 percent of smoke $PM_{2.5}$ is experienced in a different county, whereas 60 percent comes from fires in other states (Wen et al. 2023).

Figures 1 and 2 illustrate trends in days with high wildfire smoke $PM_{2.5}$ using data from Childs et al. (2022). This measure of smoke-specific $PM_{2.5}$ is in addition to any ambient $PM_{2.5}$ from traditional sources. Figure 1 plots the number of days per year that a county, weighted by population, experienced smoke-specific $PM_{2.5}$ greater than $15 \mu\text{g}/\text{m}^3$, the World Health Organization (WHO) threshold for 24-hour exposure to $PM_{2.5}$. Figure 2 shows regional trends for the WHO threshold of $15 \mu\text{g}/\text{m}^3$ and the US National Ambient Air Quality Standards 24-hour standard for $PM_{2.5}$ of $35 \mu\text{g}/\text{m}^3$ set by the EPA. The largest increases in

²These figures combine data from BLS (2019) with the analysis by Cox-Ganser and Henneberger (2021), which shows the proportions of indoor and outdoor workers by major occupations.




Figure 1 County days with smoke $PM_{2.5} > 15 \mu\text{g}/\text{m}^3$. Data from Childs et al. (2022).

extreme smoke days have been in the Pacific Northwest, Northern Rockies, and California; however, these data do not show more recent events, such as when the eastern United States was affected by large fires across Canada in the summer of 2023.

Wildfire smoke is expected to further degrade air quality as climate change exacerbates wildfire activity in the United States. Hurteau et al. (2014) estimate that wildfire emissions will increase in California by 19–101 percent by 2100. Similarly, Liu et al. (2021) use a combined climate, vegetation, and fire model to find that smoke emissions will increase by 50 percent in 2050 from the 2000 levels. Burke et al. (2023) also predict large smoke increases based




Figure 2 Regional trends in smoke days with high $PM_{2.5}$ experienced by an average resident. Data from Childs et al. (2022). Regions included in the figure are Northwest (OR, WA); Northern Rockies (ID, MT, WY); Midwest (IA, IL, IN, KS, MI, MN, MO, ND, NE, OH, SD, WI); Northeast (CT, DC, DE, MA, MD, ME, NH, NJ, NY, PA, RI, VT); California (CA); Southwest (AZ, CO, NM, NV, UT); South (OK, TX); and Southeast (AL, AR, FL, GA, KY, LA, MS, NC, SC, TN, VA, WV).

on a projected rise in vapor pressure deficit, a measure of moisture in the air that is highly correlated with wildfire activity.

Policy Responses

The two primary strategies available to policy makers and managers to reduce smoke impacts are to reduce wildfire hazard and to encourage people to avoid exposure. Whereas air quality regulation is generally the EPA's responsibility, wildfire hazard mitigation falls primarily on public land management agencies and private landowners. Land management strategies to reduce wildfire hazard include mechanical treatments, such as thinning and biomass removal, and prescribed burning to reduce dry vegetation. Especially when used together, thinning and prescribed burns can significantly reduce the severity of wildfires, provide increased opportunities for fire containment, and restore fire-adapted ecosystems to historic conditions (Schultz, McCaffrey, and Huber-Stearns 2019). Prescribed burning is seen as a particularly effective and cost-efficient way to reduce fuels in fire-adapted ecosystems in the United States (Kolden 2019). However, as a result of decades of fire exclusion and fuel buildup, fuel treatment needs in the United States are vast. In 2022, the US Forest Service (USFS) estimated that 50 million acres of public and private land in the western United States—approximately the area of Nebraska—needed to be restored over the next 10 years to reduce wildfire hazard in the highest-risk areas.

Although recent policy has increased funding for fuel treatments, including for prescribed fire, managing smoke impacts through hazard mitigation nevertheless faces significant challenges. First, capacity and funding are significant barriers to increasing application of prescribed burns (Schultz, McCaffrey, and Huber-Stearns 2019). In 2021, Clavet et al. estimated that, over the next 10 years, five to six billion dollars annually would be needed to achieve USFS targets. The Infrastructure Investment and Jobs Act (IIJA) of 2021 sizably increased the level of federal fuel treatment funding, providing \$5.5 billion for wildfire risk reduction and ecosystem restoration over the fiscal years 2022–2026, an average of \$1.1 billion per year. However, this sum will likely be insufficient to meet USFS goals. Even with further funding, the size of the current wildfire management workforce may limit the ability to dramatically increase the pace and scale of fuels management, at least in the short term. Second, agencies have generally not made smoke impacts a primary criterion for determination of priority fuel treatment locations.³ Third, on private lands, investment in fuels management is limited by private incentives, which may not take into account the full social benefit of these activities (see, e.g., Busby et al. 2012). Finally, wildfire hazard reduction may be impeded by narrow burn windows—periods of time when prescribed fire is allowable—because of the risk of escaped fires (Schultz, McCaffrey, and Huber-Stearns 2019). Combined, these factors incentivize land managers to rely on wildfires (including managed wildfires) to achieve management goals. However, wildfires are frequently larger than prescribed burns, and they may burn with greater intensity and emit more smoke.

³For example, priority “fresheds” for initial IIJA landscape investments were identified based on three criteria: potential to reduce fire risk, improved investment in underserved communities, and leveraged community partnerships (USFS 2022). Potential to reduce smoke impacts was not a primary consideration.

The fact that prescribed fires, which are intended to reduce fire hazard, can affect air quality is known as the “smoke paradox” (Schweizer and Cisneros 2017; Jones et al. 2022). This trade-off presents challenges for US air quality regulation. The Clean Air Act (CAA) functions by penalizing regions when they fail to meet air quality standards. To avoid undue punishment, the EPA treats wildfire smoke as an uncontrollable “exceptional event” and exempts smoke events from determinations of air quality attainment; a similar framework is used in Canada and Australia (Hyde et al. 2017). However, exempting smoke from prescribed burns is more difficult (Williams 2021). These rules implicitly wager that the benefits of reduced smoke from future wildfires are unlikely to outweigh the damages of certain smoke from prescribed fires. Whether this is true is an empirical question on which there is little available research. If it is not true, the CAA may need to acknowledge that wildfire hazard can be managed—in other words, that wildfires are not uncontrollable acts of God—and to use the framework of the CAA to incentivize appropriate forest management (Williams 2021).

Given the current regulatory framework, though, the EPA’s role in mitigating wildfire smoke damages has been limited to nonregulatory approaches, such as providing information about smoke impacts to encourage avoidance behaviors. For example, the EPA helps communities plan for smoke events by establishing clean air centers and caches of reserve home air filters. The agency has also partnered with the USFS to develop an online tool that provides real-time information on fire locations and air quality (GAO 2023). However, direct evidence of the public health benefits from these nonregulatory activities is lacking, and some findings indicate that benefits may be small. Treves et al. (2023) find that clean air centers are underutilized. Although related studies (e.g., Neidell 2009) find that air quality warnings can substantially increase avoidance behavior, evidence on smoke infiltration suggests that even those who avoid outdoor activity will experience large increases in indoor $PM_{2.5}$ exposure. Because of this, as well as evidence that lower-income neighborhoods engage less in protective measures than wealthy neighborhoods, Burke et al. (2022) argue that policies aimed at encouraging self-protection are likely to yield unequal benefits and to be insufficient on their own.

Conclusions

As large wildfires grow more frequent, the United States has seen increasing impacts of smoke. During extreme events, smoke increases pollution above federal attainment standards for particulate matter pollution. Smoke increases are most severe in the western states, especially California, the Pacific Northwest, and the Northern Rockies, but have affected the entire continental United States. These impacts are projected to grow over the century as the climate warms.

Both economics and social sciences literature have documented numerous impacts of wildfire smoke on human health and economic activity. These impacts are borne unequally. They include increases in mortality and severe illness, declines in maternal and infant health, and reductions in labor productivity and perceived well-being. Although much remains to be learned about damages from smoke—especially about its longer-term impacts and differences between its impacts and those of other sources of $PM_{2.5}$ —existing evidence suggests these damages are potentially even larger than direct damages from wildfires.

Current federal policy is likely insufficient to address the issue of wildfire smoke. Land management to reduce wildfire hazard faces significant challenges, including a lack of funding and regulatory hurdles for both agencies and private landowners. Air quality regulations likely are impediments to greater use of prescribed fire but could be leveraged to incentivize wildfire hazard mitigation. As much of the literature has highlighted, reliance on private adaptation alone is unlikely to adequately mitigate the damages of wildfire smoke.

References

- Aguilera, R., T. Corringham, A. Gershunov, and T. Benmarhnia. 2021. Wildfire smoke impacts respiratory health more than fine particles from other sources: Observational evidence from Southern California. *Nature Communications* 12 (1): 1493.
- Aragon, F. M., J. J. Miranda, and P. Oliva. 2017. Particulate matter and labor supply: The role of caregiving and non-linearities. *Journal of Environmental Economics and Management* 86: 295–309.
- Archsmith, J., A. Heyes, and S. Saberian. 2018. Air quality and error quantity: Pollution and performance in a high-skilled, quality-focused occupation. *Journal of the Association of Environmental and Resource Economists* 5 (4): 827–63.
- Bedi, A. S., M. Y. Nakaguma, B. J. Restrepo, and M. Rieger. 2021. Particle pollution and cognition: Evidence from sensitive cognitive tests in Brazil. *Journal of the Association of Environmental and Resource Economists* 8 (3): 443–74.
- Behrer, A. P., and S. Wang. 2022. Current benefits of wildfire smoke for yields in the US Midwest may dissipate by 2050. Policy Research Working Paper Series 9953, World Bank, Washington, DC.
- BLS (Bureau of Labor Statistics). 2019. May 2019 national occupational employment and wage estimates. BLS, Washington, DC.
- Borgschulte, M., D. Molitor, and E. Y. Zou. 2024. Air pollution and the labor market: Evidence from wildfire smoke. *Review of Economics and Statistics* 106 (6): 1558–75.
- Burke, M., M. L. Childs, B. De la Cuesta, M. Qiu, J. Li, C. F. Gould, S. Heft-Neal, and M. Wara. 2023. Wildfire influence on recent US pollution trends. NBER Working Paper no. 30882, National Bureau of Economic Research, Cambridge, MA.
- Burke, M., A. Driscoll, S. Heft-Neal, J. Xue, J. Burney, and M. Wara. 2021. The changing risk and burden of wildfire in the United States. *Proceedings of the National Academy of Sciences* 118 (2): e2011048118.
- Burke, M., S. Heft-Neal, J. Li, A. Driscoll, P. Baylis, M. Stigler, J. A. Weill, et al. 2022. Exposures and behavioural responses to wildfire smoke. *Nature Human Behaviour* 6 (10): 1351–61.
- Burkhardt, J., J. Bayham, A. Wilson, E. Carter, J. D. Berman, K. O'Dell, B. Ford, E. V. Fischer, and J. R. Pierce. 2019. The effect of pollution on crime: Evidence from data on particulate matter and ozone. *Journal of Environmental Economics and Management* 98: 102267.
- Busby, G. M., H. J. Albers, and C. A. Montgomery. 2012. Wildfire risk management in a landscape with fragmented ownership and spatial interactions. *Land Economics* 88 (3): 496–517.
- Cascio, W. E. 2018. Wildland fire smoke and human health. *Science of the Total Environment* 624: 586–95.
- Chang, T. Y., J. Graff Zivin, T. Gross, and M. Neidell. 2016. Particulate pollution and the productivity of pear packers. *American Economic Journal: Economic Policy* 8 (3): 141–69.
- . 2019. The effect of pollution on worker productivity: Evidence from call center workers in China. *American Economic Journal: Applied Economics* 11 (1): 151–72.
- Chay, K. Y., and M. Greenstone. 2005. Does air quality matter? Evidence from the housing market. *Journal of Political Economy* 113 (2): 376–424.

- Chen, S., C. Guo, and X. Huang. 2018. Air pollution, student health, and school absences: Evidence from China. *Journal of Environmental Economics and Management* 92: 465–97.
- Chen, S., P. Oliva, and P. Zhang. 2022. The effect of air pollution on migration: Evidence from China. *Journal of Development Economics* 156: 102833.
- Childs, M. L., J. Li, J. Wen, S. Heft-Neal, A. Driscoll, S. Wang, C. F. Gould, M. Qiu, J. Burney, and M. Burke. 2022. Daily local-level estimates of ambient wildfire smoke PM_{2.5} for the contiguous US. *Environmental Science and Technology* 56 (19): 13607–21.
- Clavet, C., C. Topik, M. Harrell, P. Holmes, R. Healy, and D. Wear. 2021. *Wildfire resilience funding: Building blocks for a paradigm shift*. The Nature Conservancy, Arlington, VA.
- Cox-Ganser, J. M., and P. K. Henneberger. 2021. Occupations by proximity and indoor/outdoor work: Relevance to COVID-19 in all workers and Black/Hispanic workers. *American Journal of Preventive Medicine* 60 (5): 621–28.
- Du, R., A. Mino, J. Wang, and S. Zheng. 2024. Transboundary vegetation fire smoke and expressed sentiment: Evidence from Twitter. *Journal of Environmental Economics and Management* 124: 102928.
- Fan, M., and C. Grainger. 2023. The impact of air pollution on labor supply in China. *Sustainability* 15 (17): 13082.
- Feng, S., D. Gao, F. Liao, F. Zhou, and X. Wang. 2016. The health effects of ambient PM_{2.5} and potential mechanisms. *Ecotoxicology and Environmental Safety* 128: 67–74.
- Freeman, R., W. Liang, R. Song, and C. Timmins. 2019. Willingness to pay for clean air in China. *Journal of Environmental Economics and Management* 94: 188–216.
- GAO (Government Accountability Office). 2023. Wildfire smoke: Opportunities to strengthen federal efforts to manage growing risks. GAO 23-104723, US Government Printing Office, Washington, DC.
- Gellman, J., M. Walls, and M. Wibbenmeyer. 2023. Welfare losses from wildfire smoke: Evidence from daily outdoor recreation data. RFF Working Paper Series 23-21, Resources for the Future, Washington, DC.
- Graff Zivin, J., and M. Neidell. 2012. The impact of pollution on worker productivity. *American Economic Review* 102 (7): 3652–73.
- Hamilton, T. L., and D. J. Phaneuf. 2015. An integrated model of regional and local residential sorting with application to air quality. *Journal of Environmental Economics and Management* 74: 71–93.
- He, J., H. Liu, and A. Salvo. 2019. Severe air pollution and labor productivity: Evidence from industrial towns in China. *American Economic Journal: Applied Economics* 11 (1): 173–201.
- Heft-Neal, S., A. Driscoll, W. Yang, G. Shaw, and M. Burke. 2022. Associations between wildfire smoke exposure during pregnancy and risk of preterm birth in California. *Environmental Research* 203: 111872.
- Heft-Neal, S., C. F. Gould, M. Childs, M. V. Kiang, K. Nadeau, M. Duggan, E. Bendavid, and M. Burke. 2023. Behavior mediates the health effects of extreme wildfire smoke events. NBER Working Paper no. 30969, National Bureau of Economic Research, Cambridge, MA.
- Huang, Z., and M. Skidmore. 2024. The impact of wildfires and wildfire-induced air pollution on house prices in the United States. *Land Economics* 100 (1): 22–50.
- Hurteau, M. D., A. L. Westerling, C. Wiedinmyer, and B. P. Bryant. 2014. Projected effects of climate and development on California wildfire emissions through 2100. *Environmental Science and Technology* 48 (4): 2298–304.
- Hyde, J. C., K. M. Yedinak, A. F. Talhelm, M. S. Smith, D. M. Bowman, F. H. Johnston, P. Lahm, M. Fitch, and W. T. Tinkham. 2017. Air quality policy and fire management responses addressing smoke from wildland fires in the United States and Australia. *International Journal of Wildland Fire* 26 (5): 347–63.
- Ito, K., and S. Zhang. 2020. Willingness to pay for clean air: Evidence from air purifier markets in China. *Journal of Political Economy* 128 (5): 1627–72.

- Jerrett, M., A. S. Jina, and M. E. Marlier. 2022. Up in smoke: California's greenhouse gas reductions could be wiped out by 2020 wildfires. *Environmental Pollution* 310: 119888.
- Johnston, F. H., S. B. Henderson, Y. Chen, J. T. Randerson, M. Marlier, R. S. DeFries, P. Kinney, D. M. Bowman, and M. Brauer. 2012. Estimated global mortality attributable to smoke from landscape fires. *Environmental Health Perspectives* 120 (5): 695–701.
- Jolly, W. M., M. A. Cochrane, P. H. Freeborn, Z. A. Holden, T. J. Brown, G. J. Williamson, and D. M. Bowman. 2015. Climate-induced variations in global wildfire danger from 1979 to 2013. *Nature Communications* 6 (1): 7537.
- Jones, B. A. 2017. Are we underestimating the economic costs of wildfire smoke? An investigation using the life satisfaction approach. *Journal of Forest Economics* 27: 80–90.
- Jones, B. A., S. McDermott, P. A. Champ, and R. P. Berrens. 2022. More smoke today for less smoke tomorrow? We need to better understand the public health benefits and costs of prescribed fire. *International Journal of Wildland Fire* 31 (10): 918–26.
- Kim, K., E. Kabir, and S. Kabir. 2015. A review on the human health impact of airborne particulate matter. *Environment International* 74: 136–43.
- Kolden, C. A. 2019. We're not doing enough prescribed fire in the western United States to mitigate wildfire risk. *Fire* 2 (2): 30.
- Liang, Y., D. Sengupta, M. J. Campmier, D. M. Lunderberg, J. S. Apte, and A. H. Goldstein. 2021. Wildfire smoke impacts on indoor air quality assessed using crowdsourced data in California. *Proceedings of the National Academy of Sciences* 118 (36): e2106478118.
- Liu, J. C., and R. D. Peng. 2019. The impact of wildfire smoke on compositions of fine particulate matter by ecoregion in the western US. *Journal of Exposure Science and Environmental Epidemiology* 29 (6): 765–76.
- Liu, J. C., G. Pereira, S. A. Uhl, M. A. Bravo, and M. L. Bell. 2015. A systematic review of the physical health impacts from non-occupational exposure to wildfire smoke. *Environmental Research* 136: 120–32.
- Liu, Y., Y. Liu, J. Fu, C.-E. Yang, X. Dong, H. Tian, B. Tao, et al. 2021. Projection of future wildfire emissions in western USA under climate change: Contributions from changes in wildfire, fuel loading and fuel moisture. *International Journal of Wildland Fire* 31 (1): 1–13.
- Marlier, M. E., K. I. Brenner, J. C. Liu, L. J. Mickley, S. Raby, E. James, R. Ahmadov, and H. Riden. 2022. Exposure of agricultural workers in California to wildfire smoke under past and future climate conditions. *Environmental Research Letters* 17 (9): 094045.
- Marlon, J. R., P. J. Bartlein, D. G. Gavin, C. J. Long, R. S. Anderson, C. E. Briles, K. J. Brown, et al. 2012. Long-term perspective on wildfires in the western USA. *Proceedings of the National Academy of Sciences* 109 (9): E535–E543.
- McCoy, S. J., and X. Zhao. 2021. Wildfire and infant health: A geospatial approach to estimating the health impacts of wildfire smoke exposure. *Applied Economics Letters* 28 (1): 32–37.
- Nam, K. M., Y. Ou, E. Kim, and S. Zheng. 2022. Air pollution and housing values in Korea: A hedonic analysis with long-range transboundary pollution as an instrument. *Environmental and Resource Economics* 82 (2): 383–407.
- Neidell, M. 2009. Information, avoidance behavior, and health: The effect of ozone on asthma hospitalizations. *Journal of Human Resources* 44 (2): 450–78.
- O'Dell, K., B. Ford, J. Burkhardt, S. Magzamen, S. C. Anenberg, J. Bayham, E. V. Fischer, and J. R. Pierce. 2022. Outside in: The relationship between indoor and outdoor particulate air quality during wildfire smoke events in western US cities. *Environmental Research: Health* 1 (1): 015003.
- O'Dell, K., B. Ford, E. V. Fischer, and J. R. Pierce. 2019. Contribution of wildland-fire smoke to US PM_{2.5} and its influence on recent trends. *Environmental Science and Technology* 53 (4): 1797–804.

- Power, M. C., M. A. Kioumourtzoglou, J. E. Hart, O. I. Okereke, F. Laden, and M. G. Weisskopf. 2015. The relation between past exposure to fine particulate air pollution and prevalent anxiety: Observational cohort study. *British Medical Journal* 350: h1111.
- Qiu, M., J. Li, C. F. Gould, R. Jing, M. Kelp, M. Childs, M. Kiang, S. Heft-Neal, N. Diffenbaugh, and M. Burke. 2024. Mortality burden from wildfire smoke under climate change. NBER Working Paper no. 32307, National Bureau of Economic Research, Cambridge, MA.
- Reid, C. E., M. Brauer, F. H. Johnston, M. Jerrett, J. R. Balmes, and C. T. Elliott. 2016. Critical review of health impacts of wildfire smoke exposure. *Environmental Health Perspectives* 124 (9): 1334–43.
- Richardson, L. A., P. A. Champ, and J. B. Loomis. 2012. The hidden cost of wildfires: Economic valuation of health effects of wildfire smoke exposure in Southern California. *Journal of Forest Economics* 18 (1): 14–35.
- Roberts, G., and M. J. Wooster. 2021. Global impact of landscape fire emissions on surface level PM_{2.5} concentrations, air quality exposure and population mortality. *Atmospheric Environment* 252: 118210.
- Rubin, N. B., and G. Wong-Parodi. 2022. As California burns: The psychology of wildfire- and wildfire smoke-related migration intentions. *Population and Environment* 44 (1–2): 15–45.
- Ruminski, M., J. Simko, J. Kibler, S. Kondragunta, R. Draxler, P. Davidson, and P. Li. 2008. Use of multiple satellite sensors in NOAA's operational near real-time fire and smoke detection and characterization program. *Proceedings of SPIE 7089, Remote Sensing of Fire: Science and Application* 70890A: 76–86.
- Sass, V., N. Kravitz-Wirtz, S. M. Karceski, A. Hajat, K. Crowder, and D. Takeuchi. 2017. The effects of air pollution on individual psychological distress. *Health and Place* 48: 72–79.
- Schmidt, C. W. 2022. Well played: Using game app data to assess wildfire smoke and cognitive performance. *Environmental Health Perspectives* 130 (7): 074002.
- Schultz, C. A., S. M. McCaffrey, and H. R. Huber-Stearns. 2019. Policy barriers and opportunities for prescribed fire application in the western United States. *International Journal of Wildland Fire* 28 (11): 874–84.
- Schweizer, D. W., and R. Cisneros. 2017. Forest fire policy: Change conventional thinking of smoke management to prioritize long-term air quality and public health. *Air Quality Atmosphere and Health* 10: 33–36.
- Senande-Rivera, M., D. Insua-Costa, and G. Miguez-Macho. 2022. Spatial and temporal expansion of global wildland fire activity in response to climate change. *Nature Communications* 13 (1): 1208.
- Treves, R. J., E. Liu, S. L. Fischer, E. Rodriguez, and G. Wong-Parodi. 2023. Wildfire smoke clean air centers: Identifying barriers and opportunities for improvement from California practitioner and community perspectives. *Society and Natural Resources* 36 (9): 1078–97.
- Urbanski, S. P., W. M. Hao, and S. Baker. 2008. Chemical composition of wildland fire emissions. *Developments in Environmental Science* 8: 79–107.
- USFS (US Forest Service). 2022. Confronting the wildfire crisis: Initial landscape investments to protect communities and improve resilience in America's forests. No. FS-1187d, USFS, Washington, DC.
- Wang, D., D. Guan, S. Zhu, M. M. Kinnon, G. Geng, Q. Zhang, H. Zheng, et al. 2021. Economic footprint of California wildfires in 2018. *Nature Sustainability* 4 (3): 252–60.
- Wegesser, T. C., K. E. Pinkerton, and J. A. Last. 2009. California wildfires of 2008: Coarse and fine particulate matter toxicity. *Environmental Health Perspectives* 117 (6): 893–97.
- Wen, J., and M. Burke. 2022. Lower test scores from wildfire smoke exposure. *Nature Sustainability* 5 (11): 947–55.

- Wen, J., S. Heft-Neal, P. Baylis, J. Boomhower, and M. Burke. 2023. Quantifying fire-specific smoke exposure and health impacts. *Proceedings of the National Academy of Sciences* 120 (51): e2309325120.
- Whiting, J., and M. Krstic. 2007. Impact of bushfire smoke on grape and wine. Study report, Department of Primary Industries, Victoria, Australia.
- Williams, E. 2021. Reimagining exceptional events: Regulating wildfires through the Clean Air Act. *Washington Law Review* 96: 765–810.